## K<sub>S</sub><sup>0</sup> production in Pb+Pb collisions at CERN SPS

S. Margetis and the NA49 Collaboration

Neutral kaons were reconstructed in NA49 Time Projection Chambers (TPC) using a method described in [1], via their V0 decay topology. A sample of 20000 Pb+Pb events at 158 GeV/c was analyzed and 15000 simulated events were used for efficiency and background estimations. The interaction cross-section is 5% of  $\sigma_{inel}^{tot}$  which corresponds to b  $_{max}=3$  fm. Results were obtained in the rapidity region 2.0 < y < 2.8 and  $p_T > 0.4 \,\mathrm{GeV/c}$ .

The resulting transverse mass distribution is shown in Fig. 1. An exponential fit  $(1/m_T dN/dm_T = e^{-m_T/T})$  gives an inverse slope parameter (T) of  $220 \pm 15$  MeV.




Figure 1: Transverse mass distributions of K  $_{S}^{0}$  in the rapidity region 2.2 < y < 2.6.

By integrating over all  $p_T$  values one obtains the rapidity density (dN/dy) which is shown in Fig. 2. The filled symbols are the measured data whereas the open ones are the same values but reflected around mid-rapidity  $(y_{mid}^{Pb+Pb} = 2.92)$ . This is because Pb+Pb is a symmetric system and assuming the trigger biases are negligible. The solid line is the fitted S+S distribution scaled with the ratio of number of participants in the two systems (390/58 = 6.7) which

is also the ratio of produced negative hadrons (mostly pions). The two distributions are compatible within errors. This leads to the conclusion that the  $K_S^0/\pi$  ratio is about the same in S+S and Pb+Pb collisions [2]. This ratio in S+S was found to be twice as large than in nucleonnucleon collisions at the same energy [3]. It is therefore intriguing that in a much larger system, like Pb+Pb, there is no further enhancement of this ratio.




Figure 2: Rapidity distribution of K  $_{S}^{0}$  in Pb+Pb collisions (symbols). See text for an explanation of the symbols.

## References

- [1] S. Margetis, Elsewhere in this report
- [2] S. Margetis, Heavy Ion Phys. 4 (96) 63
- [3] T. Alber et al, NA35 Collaboration, Z.Phys.C 64(94)195